arXiv:1105.1154v3 [math.GT] 17 Sep 2012

Fuchsian subgroups of lattices acting on hermitian symmetric spaces

Ted Chinburg[*]
University of Pennsylvania
ted@math.upenn.edu

Matthew Stover[†]
University of Michigan
stoverm@umich.edu

January 20, 2013

1 Introduction

Let S be a smooth compact arithmetic quotient of a two-dimensional hermitian symmetric domain, and suppose that S contains an arithmetic Fuchsian curve, i.e., a totally geodesic holomorphic immersion of a smooth projective curve of genus $g \geq 2$. In this paper, we will show that there is a connected divisor D on S with all irreducible components Fuchsian curves such that the fundamental group of D generates a subgroup of finite index in the fundamental group of S. This is an example of the principle that one can understand the geometry of higher dimensional manifolds via their totally geodesic submanifolds; for other examples see [25, 6, 23, 19, 7, 2, 27].

One consequence of our results is the following structure theorem for the Albanese varieties of arithmetic complex hyperbolic 2-manifolds.

Theorem 1.1. *Let* $\Gamma < \mathrm{SU}(2,1)$ *be a torsion free cocompact arithmetic lattice of simple type (see §2.3). There exists* $r \geq 1$ *and Fuchsian curves* $C_1, \dots, C_r$ *on* $S = \Gamma \backslash \mathbf{H}^2_{\mathbb{C}}$ *such that if the Albanese variety* $\mathrm{Alb}(S)$ *is nontrivial, then every simple factor of* $\mathrm{Alb}(S)$ *is isogenous to a factor of the Jacobian of the normalization* $C_j^{\#}$ *of (at least) one of the curves* C_j.

Here $C_j^{\#}$ is the quotient of the upper half plane by the maximal Fuchsian subgroup of Γ determined by the totally geodesic curve C_j.

---

[*] partially supported by NSF grant DMS 0801030

[†] partially supported by NSF RTG grant DMS 0602191

If Γ is a congruence arithmetic lattice of simple type, Gelbart and Rogawski [8] proved that the first cohomology group of $\Gamma\backslash\mathbf{H}^2_{\mathbb{C}}$, which determines $\mathrm{Alb}(\Gamma\backslash\mathbf{H}^2_{\mathbb{C}})$, arises from the theta correspondence. Murty and Ramakrishnan then used Falting's work on the Mordell conjecture in [20] to show that the simple factors of $\mathrm{Alb}(\Gamma\backslash\mathbf{H}^2_{\mathbb{C}})$ are, in fact, CM abelian varieties. This gave a positive answer to a question of Langlands.

Theorem 1.1 provides information of a different nature about $\mathrm{Alb}(\Gamma\backslash\mathbf{H}^2_{\mathbb{C}})$ for both congruence and noncongruence Γ, namely that $\mathrm{Alb}(\Gamma\backslash\mathbf{H}^2_{\mathbb{C}})$ is built from the Jacobians of Fuchsian curves on $\Gamma\backslash X$. By work of Kazhdan in [14], one can always find elements in the commensurability class with nontrivial Albanese variety, and there are always noncongruence groups in the commensurability classes under consideration. It would be interesting to know whether or not the factors which appear for noncongruence Γ must also have complex multiplication.

For a divisor $D = \sum \alpha_j C_j$, where $\alpha_j > 0$ and the C_j are distinct and irreducible, let $|D|$ be the compact analytic subspace of the complex points of S determined by the union of the spaces defined by the C_j. We now give a precise statement of our main result.

Theorem 1.2. *Let G be $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ or $\mathrm{SU}(2,1)$ with corresponding hermitian symmetric space X, and let Γ be a cocompact arithmetic lattice in G. Assume that the complex algebraic surface $S = \Gamma\backslash X$ contains a holomorphically immersed totally geodesic projective algebraic curve. Then there exist finitely many such curves $C_1, \dots, C_r \subset S$ and positive integers $\alpha_1, \dots, \alpha_r$ such that:*

1. *The divisor $D = \sum \alpha_j C_j$ is a connected effective divisor on S.*
2. *The image of $\pi_1(|D|)$ in Γ under the natural homomorphism is a finite index subgroup.*

See the remark at the end of §3 for a discussion of how one can bound the number of curves in Theorem 1.2 from above. The arithmetic assumption is actually superfluous for $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$. Indeed, nonarithmetic lattices are reducible by Margulis's Arithmeticity Theorem [17], and Theorem 1.2 is easy to prove for reducible lattices, which contain a direct product of two Fuchsian groups with finite index. Therefore, we implicitly assume throughout this paper that lattices in $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ are irreducible.

We now outline the rest of the paper. In §2, we describe the arithmetic of the lattices under consideration. Important for us is the existence of infinitely many commensurability classes of Fuchsian curves on the arithmetic surfaces under consideration. Since it does not seem to appear in the literature, we give an explicit parameterization of Fuchsian subgroups of arithmetic lattices in $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ and $\mathrm{SU}(2,1)$ associated with holomorphically immersed totally geodesic curves. In both cases, we give our parameterization in terms of the well-known description of arithmetic Fuchsian groups via quaternion algebras; see e.g. [28, 16].

We prove Theorem 1.2 in §3, where we construct a collection of Fuchsian curves that allows us to apply a Weak Lefschetz Theorem. Weak Lefschetz Theorems go back to Nori's work on Zariski's conjecture [22], but the version we apply is due to Napier and Ramachandran [21]. See also [5]. Our application to Albanese varieties of complex hyperbolic manifolds is proven in §4.

Acknowledgments

The first author thanks J. B. Bost for a very helpful conversation about Lefschetz Theorems. The second author thanks Robert Lazarsfeld for conversation related to §3, the University of Pennsylvania for its hospitality, and Alan Reid for conversations related to this project and for his part in instigating this collaboration. Both authors thank Misha Kapovich for noticing an error in an early draft of this paper and Domingo Toledo for helpful comments.

2 Arithmetic of lattices acting on hermitian symmetric spaces

2.1 Generalities on lattices

Let G be a semisimple Lie group with finite center and associated symmetric space X of noncompact type. Then X is called *Hermitian symmetric* when it carries a G-invariant complex structure [13, Chapter VIII]. The quotient of X by any cocompact lattice Γ in G is a projective algebraic variety [1], and $\Gamma\backslash X$ is a smooth variety when Γ is torsion free. Suppose H is a semisimple Lie subgroup of G with finite center and associated symmetric space Y. There is then a totally geodesic embedding $Y \to X$, and the closed totally geodesic subspaces of $\Gamma\backslash X$ arise from the subgroups H of this kind such that $\Gamma \cap H$ is a lattice in H; see [30, §1.4].

Now suppose that K_0 is a number field and that $\mathcal{G}$ is a K_0-algebraic group such that $\mathcal{G}(K_0 \otimes_{\mathbb{Q}} \mathbb{R})$ modulo compact factors is isomorphic to G (see [30, §3.6] for definitions). Let Γ be an arithmetic lattice commensurable with the lattice of K_0-integral points of $\mathcal{G}$. Then connected totally geodesic submanifolds of $\Gamma\backslash X$ are determined by the semisimple F-algebraic subgroups $\mathcal{H}$ of $\mathcal{G}$ for some field $F \subseteq K_0$, and $\Gamma \cap \mathcal{H}(\mathbb{R})$ is an arithmetic lattice in $\mathcal{H}(\mathbb{R})$.

The cases of interest here are where G is either $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ or $\mathrm{SU}(2,1)$, and we recall the construction of arithmetic lattices in $\mathrm{SL}_2(\mathbb{R})$ and $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ in §2.2 and consider $\mathrm{SU}(2,1)$ in §2.3. The respective hermitian symmetric domains are the product $\mathbf{H}^2 \times \mathbf{H}^2$ of two hyperbolic planes or the complex hyperbolic plane $\mathbf{H}^2_{\mathbb{C}}$.

The possible holomorphically embedded totally geodesic curves are *projective Fuchsian curves*, that is, quotients of $\mathbf{H}^2$ by cocompact lattices in $\mathrm{SL}_2(\mathbb{R})$. Note that the holomorphic isometry group $\mathrm{Isom}_h(\mathbf{H}^2)$ of $\mathbf{H}^2$ is naturally isomorphic to $\mathrm{PSL}_2(\mathbb{R}) \cong \mathrm{PU}(1,1)$. By a *cocompact Fuchsian group*, we will mean a discrete subgroup of any of the groups $\mathrm{SL}_2(\mathbb{R})$, $\mathrm{PSL}_2(\mathbb{R})$, $\mathrm{U}(1,1)$, $\mathrm{SU}(1,1)$, or $\mathrm{PU}(1,1)$ such that the image in $\mathrm{Isom}_h(\mathbf{H}^2)$ is a cocompact lattice. See [28] and [16] for an account of the basic theory of arithmetic Fuchsian groups. We also give a complete classification of the possible arithmetic Fuchsian subgroups determined by holomorphically embedded totally geodesic projective curves on compact arithmetic quotients of $\mathbf{H}^2 \times \mathbf{H}^2$ and $\mathbf{H}^2_{\mathbb{C}}$ in §§2.2-2.3; these classifications to not appear to be in the literature.

2.2 Lattices in $\mathrm{SL}_2(\mathbb{R})^n$ acting on $(\mathbf{H}^2)^n$ for $1 \leq n \leq 2$

We recall from [4, §3] the construction of irreducible arithmetic lattices acting on products of hyperbolic planes. Let $\mathbf{H}^2$ be the hyperbolic plane. For $n = 1, 2$ let G (resp. X) be the product of n copies of $\mathrm{SL}_2(\mathbb{R})$ (resp. $\mathbf{H}^2$). Let K be a totally real number field and A a quaternion algebra over K. We assume that A is unramified at exactly n archimedean places of K. Then there is an algebra surjection $\pi : \mathbb{R} \otimes_K A \to \mathrm{M}_2(\mathbb{R})^n$ whose kernel is a product of copies of the Hamilton quaternion algebra over $\mathbb{R}$.

Let $\mathcal{O}$ be an order of A and let $\mathcal{O}^1$ be the multiplicative subgroup of elements with reduced norm one to K. Then $\pi(\mathcal{O}^1)$ is a lattice in $\mathrm{SL}_2(\mathbb{R})^n$. A lattice $\Gamma < \mathrm{SL}_2(\mathbb{R})^n$ is arithmetic when a conjugate of Γ is commensurable with a group $\pi(\mathcal{O}^1)$ constructed in the above way. We will always replace Γ with a conjugate and assume that it is commensurable with $\pi(\mathcal{O}^1)$.

When $n = 1$ such Γ are arithmetic Fuchsian groups. When $n = 2$, a lattice $\Gamma < G$ is *irreducible* if its projection onto any factor is dense in the analytic topology. A lattice fails to be irreducible in this case if and only if there is a finite index subgroup that is the direct product of two Fuchsian groups. All irreducible lattices in G are arithmetic by Margulis's Arithmeticity Theorem [17].

For $n = 1, 2$, two arithmetic lattices Γ_1 and Γ_2 are commensurable if and only if the associated quaternion algebras are isomorphic. Therefore, up to conjugacy in $\mathrm{SL}_2(\mathbb{R})^n$, commensurability classes of irreducible arithmetic lattices acting on $(\mathbf{H}^2)^n$ are in one-to-one correspondence with totally real fields K and K-quaternion algebras ramified at all but n real places of K. An arithmetic lattice Γ is cocompact if and only if $A \neq \mathrm{M}_2(K)$.

Proposition 2.1. *Suppose $n = 2$ and $\Gamma \subset \mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$ is an irreducible lattice commensurable with $\pi(\mathcal{O}^1)$ for an order $\mathcal{O}$ in a quaternion algebra A over the totally real number field K. Then, there is a one-to-one correspondence between*

1. *commensurability classes of arithmetic Fuchsian subgroups of Γ and*
2. *degree two subfields $K_0 \subset K$ and quaternion K_0-subalgebras $B \subset A$ such that $A = K \otimes_{K_0} B$.*

In particular, Γ contains infinitely many commensurability classes of arithmetic Fuchsian groups if and only if it contains one arithmetic Fuchsian subgroup.

Proof. The correspondence between (*1*) and (*2*) follows from analyzing the possible simple subalgebras of A. The last statement follows from the fact that $K \otimes_{K_0} B = K \otimes_{K_0} B'$ whenever B' is another quaternion algebra over K_0 such that the places of K_0 that ramify in exactly one of B or B' do not split in K. $\square$

2.3 Lattices acting on $\mathbf{H}^2_{\mathbb{C}}$

First, we quickly recall the standard construction of $\mathbf{H}^2_{\mathbb{C}}$. See [9, Chapter 3] for further details. Let V be a rank 3 vector space over $\mathbb{C}$ and h a hermitian form on V of signature $(2, 1)$. If V_- denotes the subspace of h-negative vectors, then $\mathbf{H}^2_{\mathbb{C}}$ is the space $\mathbb{P}(V_-)$ of h-negative lines. There is a natural biholomorphism from $\mathbf{H}^2_{\mathbb{C}}$ with the metric determined by h to the unit ball in $\mathbb{C}^2$ with the Bergman metric.

Holomorphic embeddings of $\mathbf{H}^2$ in $\mathbf{H}^2_{\mathbb{C}}$ arise from rank 2 subspaces of V on which the restriction of h to V has signature $(1, 1)$. Taking the h-orthogonal complement, it follows that holomorphic totally geodesic embeddings of $\mathbf{H}^2$ into $\mathbf{H}^2_{\mathbb{C}}$ are in one-to-one correspondence with h-positive lines in V. See [9, §3.3.1] and [2, §12.2].

By [2, Chapter 8] the only arithmetic lattices in $\mathrm{SU}(2, 1)$ which contain totally geodesic surfaces are those of so-called *simple type* (also sometimes called first type). These are defined as follows.

Recall that a *CM pair* K/K_0 is a totally imaginary quadratic extension of a totally real number field. Let $z \mapsto \overline{z}$ be the nontrivial automorphism of K over K_0 and h be a hermitian form on K^3 such that h is indefinite at precisely one $\mathrm{Gal}(K/K_0)$ pair of embeddings of K into $\mathbb{C}$. Replacing h with $-h$ if necessary, we assume that h has signature $(2,1)$ at the place of K where it is indefinite.

If $\mathcal{O}$ is the integer ring of K, the K_0-algebraic group

$$\mathrm{SU}(h) = \{x \in \mathrm{SL}_3(K) \ : \ {}^t\overline{x}hx = h\},$$

contains the discrete subgroup $\Gamma^1_{\mathcal{O}} = \mathrm{SU}(h) \cap \mathrm{SL}_3(\mathcal{O})$. Then $\Gamma^1_{\mathcal{O}}$ projects to a lattice in $\mathrm{SU}(2,1)$, since h is indefinite above exactly one real embedding of K_0. It is known (see [24, §1.2]) that the commensurability class of $\Gamma^1_{\mathcal{O}}$ in $\mathrm{SU}(2,1)$ depends only on K, not the choice of hermitian form. The lattice $\Gamma^1_{\mathcal{O}}$ is cocompact if and only if $K_0 \neq \mathbb{Q}$.

The following classifies and parametrizes the possible commensurability classes of arithmetic Fuchsian subgroups of arithmetic lattices in $\mathrm{SU}(2,1)$ of simple type. Since we do not know of a reference in the literature we give a complete proof.

Theorem 2.2. *Let $\Gamma < \mathrm{SU}(2,1)$ be an arithmetic lattice of simple type with associated CM pair K/K_0. Then, there is a one-to-one correspondence between*

1. *commensurability classes of arithmetic Fuchsian subgroups of Γ corresponding to totally geodesic projective algebraic curves on $\Gamma\backslash\mathbf{H}^2_{\mathbb{C}}$ and*
2. *K_0-quaternion algebras A ramified at all but one infinite place of K_0 and at any finite set of nonarchimedean places which do not split in K/K_0.*

In particular, any such Γ contains infinitely many distinct commensurability classes of such arithmetic Fuchsian subgroups.

Proof. Assume that h is a hermitian form on $V = K^3$ that is indefinite above exactly one archimedean place of K_0. Let $W \cong K^2 \subset V$ be the h orthogonal complement of a line in V which is h-positive at the archimedean place of K_0 over which h is indefinite, and let h_W be the restriction of h to W. By our earlier discussion of holomorphic embeddings of $\mathbf{H}^2$ into $\mathbf{H}^2_{\mathbb{C}}$, the pair (W, h_W) determines a maximal arithmetic Fuchsian subgroup Σ_W of Γ associated to a holomorphic curve on $\Gamma\backslash\mathbf{H}^2_{\mathbb{C}}$, and all such curves arise in this way. We now proceed to determine the K_0-quaternion algebra associated to the commensurability class of this arithmetic Fuchsian group Σ_W.

After choosing a K-basis for W, there is a natural homomorphism from Σ_W to $\mathrm{GL}_2(K)$. Furthermore, the K_0-subalgebra B of $\mathrm{M}_2(K)$ generated by Σ_W is four-dimensional and noncommutative, so it is a quaternion algebra. Since h_W is indefinite at exactly one infinite place of K_0 and B embeds into $\mathrm{M}_2(K)$, B must satisfy condition (*2*) of the theorem.

Conversely, suppose B is a quaternion algebra over K_0 satisfying the conditions in (*2*). We must show that there is a commensurability class of arithmetic Fuchsian subgroups of Γ associated with B. Condition (*2*) implies that we have an embedding of K into B, so B is a two-dimensional left vector space over K. The nontrivial automorphism of K over K_0 extends to an anti-involution $z \to \overline{z}$ of B over K_0.

There is a left K-linear action of B on itself under which $z \in B$ acts on $x \in B$ by $x \mapsto x \cdot \overline{z}$. This is a K_0-algebra embedding of B into $\mathrm{Mat}_2(K)$. Let $\mathrm{Tr} : B \to K$ be the composition of this embedding with the trace. Identifying B with K^2, we obtain a hermitian form on K^2 by

$$h_B(x, y) = \mathrm{Tr}(x\overline{y}),$$

for $x, y \in B$.

Now, consider the hermitian form H_B on K^3 defined as follows. Write $K^3 = W \oplus \ell$, where W is two-dimensional over K and ℓ is a line over K. Let the restriction of H_B to W be h_B, let ℓ be H_B-orthogonal to W, and let the H_B-norm of some generator of ℓ be $\det(h)/\det(h_B)$ in K_0. Then $\det(h)/\det(h_B)$ is totally positive since condition (*2*) implies that $\det(h_B)$ has the same sign as $\det(h)$ at every archimedean place of K_0. By [26, Chapter 10, Example 1.8(iii)], K^3 equipped with H_B is isomorphic as a hermitian vector space to K^3 equipped with h. Let B^1 be the group of elements in B with reduced norm 1 to K_0. Then the action of B^1 stabilizes h_B, so B^1 embeds in the stabilizer of W inside $\mathrm{SU}(h)$. We conclude that $\Gamma \cap B$ is an arithmetic Fuchsian subgroup of Γ and that the commensurability class of this group is associated with B. This completes the proof. □

3 The proof of Theorem 1.2

In this section we assume familiarity with basic complex algebraic geometry, e.g., from [11]. Let S be a complex surface, $D = \sum \alpha_j C_j$ be a divisor on S with irreducible components C_j, and let $(\, , \,)$ denote the intersection pairing on S. Since this pairing only depends on the homology class of an algebraic cycle on S, we will abuse notation and use D to denote the divisor or its class in $H^2(S; \mathbb{Z})$. A divisor D is *positive* when $(D, D) > 0$.

We will say D is *reduced* if $D = \sum C_j$ for some finite set $\{C_j\}$ of distinct irreducible curves, and a divisor D' is *supported by* D if $D' = \sum \alpha_j C_j$ for some coefficients $\alpha_j \geq 0$. We say D' is *precisely* supported by D if $\alpha_j > 0$ for every j, i.e., if D' is an effective divisor with irreducible components exactly the irreducible components of D. For any D' supported by D, let $|D'|$ be compact analytic subspace of the complex points of S that is the union of the spaces defined by the C_j for which $\alpha_j > 0$. If D' is precisely supported by D, then $\pi_1(|D'|)$ and $\pi_1(|D|)$ are both equal, by definition, to the fundamental group of the topological subspace of S that is the union of the complex points of the C_j; note that this is not in general the subgroup of $\pi_1(S)$ generated by the groups $\pi_1(|C_j|)$.

We now introduce some notation specific to our situation. Let $S = \Gamma \backslash X$ be a quotient of either $\mathbf{H}^2 \times \mathbf{H}^2$ or $\mathbf{H}^2_{\mathbb{C}}$ by an irreducible cocompact lattice. To prove Theorem 1.2 it suffices to pass to a subgroup of finite index, so we may assume Γ is torsion free and S is smooth. We keep this assumption for the remainder of this section.

A *Fuchsian curve* on S is a totally geodesic holomorphic immersion of a cocompact quotient of $\mathbf{H}^2$ into S. Fuchsian curves are always irreducible. A *Fuchsian divisor* on S is a divisor $D = \sum \alpha_j C_j$ such that every irreducible component C_j is a Fuchsian curve, where the α_j are arbitrary integers. We call D a *reduced Fuchsian divisor* when $D = \sum C_j$ is a reduced divisor on S such that the C_j are distinct Fuchsian curves.

To prove Theorem 1.2, we apply a so-called Weak Lefschetz Theorem. The first such theorem is due to Nori [22], with later contributions by Napier–Ramachandran [21] and Campana [5]. Our approach is closely related to [5], but uses a result from [21]. Our proof uses the following result, which follows from setting $R = \emptyset$ in Theorem 1.1 of [21] (see also [5, Lemma 2.5 and Prop. A.B.1]).

Theorem 3.1 (Napier–Ramachandran)**.** *Let U be a connected complex manifold, let S be a connected smooth projective variety of dimension $n > 1$, let $\Phi : U \to S$ be a holomorphic map, let Y be a connected compact analytic subspace (not necessarily reduced) of U, and let $\widehat{U}$ be the formal completion of U with respect to Y. Assume that*

(i) Φ is locally biholomorphic, and

(ii) $\dim \mathrm{H}^0(\widehat{U}, \widehat{\mathcal{O}(\Phi^* L)}) < \infty$ for every holomorphic line bundle L on S.

Then the image of $\pi_1(Y) \to \pi_1(S)$ has finite index.

Our Weak Lefschetz Theorem for Fuchsian divisors is the following.

Theorem 3.2. *Let* $D = \sum C_j$ *be a reduced Fuchsian divisor on* $S = \Gamma\backslash X$*, where* X *is* $\mathbf{H}^2 \times \mathbf{H}^2$ *or* $\mathbf{H}^2_{\mathbb{C}}$ *and* Γ *is a torsion free cocompact lattice in* $\mathrm{Isom}(X)$*. Suppose that the support of* D *is connected and that* D *precisely supports a positive effective Fuchsian divisor* D' *on* S *such that*

$$(D', C_j) > 0 \text{ for all } j. \tag{$\star$}$$

Let $\pi_{D'} : S_{D'} \to S$ *be the étale covering of* S *determined by the image* $\Gamma_{D'}$ *of* $\pi_1(|D'|) = \pi_1(|D|)$ *in* Γ *under the homomorphism induced by the inclusion on* D' *into* S*. Then* $\pi_{D'}$ *is a finite morphism, i.e., the image of* $\pi_1(|D'|)$ *in* Γ *is a finite index subgroup.*

Proof. By definition, $S_{D'}$ is the maximal covering of S to which a sufficiently small topological neighborhood of $|D'|$ lifts isomorphically. Let $\widehat{D}$, $\widehat{D}'$, and $\widehat{C}_j$ denote the lifts of D, D', and C_j to $S_{D'}$. The analytic spaces defined by these divisors are isomorphic to their images in S under the covering projection, and $\widehat{D}'$ is a positive Fuchsian divisor on $S_{D'}$ satisfying ($\star$).

Let $Y = |\widehat{D}'|$ and let U be a small connected neighborhood of Y in $S_{D'}$. Then Y is a connected compact analytic subspace of U, though it is not necessarily reduced. We may assume that U is a complex manifold, and then the restriction $\Phi : U \to S$ of the covering map $\pi : S_{D'} \to S$ is a local biholomorphism. To apply Theorem 3.1 we must show that condition (*ii*) of this theorem holds.

Let $\widehat{U}$ be the formal completion of U with respect to Y. The ideal sheaf $I = I_Y$ of Y is $\mathcal{O}_U(-Y)$, and structure sheaf of the k^{th} infinitesimal neighborhood Y_k of Y in U is $\mathcal{O}_{Y_k} = \mathcal{O}_U/\mathcal{O}_U(-kY)$. For any line bundle L on S, we have

$$\mathrm{H}^0(\widehat{U}, \widehat{\mathcal{O}(\Phi^* L)}) = \varprojlim_k \mathrm{H}^0(Y_k, \Phi^* L \otimes \mathcal{O}_{Y_k}), \tag{1}$$

and we have an exact sequence

$$0 \to \mathrm{H}^0(Y, \Phi^* L \otimes (I^k/I^{k+1})) \to \mathrm{H}^0(Y_{k+1}, \Phi^* L \otimes \mathcal{O}_{Y_{k+1}}) \to \mathrm{H}^0(Y_k, \Phi^* L \otimes \mathcal{O}_{Y_k}).$$

Therefore to show that (1) is finite dimensional, as required in condition (*ii*) of Theorem 3.1, it suffices to show that $\mathrm{H}^0(Y, \Phi^* L \otimes (I^k/I^{k+1})) = 0$ if k is sufficiently large. Since Y is a Cartier divisor on the smooth surface S, it is Cohen–Macaulay and projective. By Serre duality (see [12, Thm. III.7.6(ii)]) together with flat base change for cohomology, it suffices to show that $(I/I^2)^{-1} = \mathcal{O}(Y)|_Y$ is an ample line bundle on Y. By [15, Example 1.2.28(ii)], it is enough to prove that for all j the Euler characteristic

$\chi(\mathcal{O}(mY)|_{\widehat{C}_j})$ goes to $+\infty$ as $m \to +\infty$. This is an immediate consequence of condition $(\star)$ together with the Riemann–Roch Theorem for the (possibly singular) curve $\widehat{C}_j$ (see [12, Ex. IV.1.9]). Therefore Theorem 3.1 applies, so the image of $\pi_1(|\widehat{D}'|) = \pi_1(|D'|)$ in $\pi_1(S)$ has finite index. This completes the proof of the theorem. $\square$

To prove Theorem 1.2, we now must show that any S satisfying the conditions of the theorem contains a divisor D as in Theorem 3.2. We begin with the following lemma (see [5, Lemma A.A.1]).

Lemma 3.3. *Let S be a complex surface, not necessarily compact, and let C be a reduced compact connected curve on S. Write $C = \bigcup C_j$ for a collection $\{C_j\}_j$ of distinct irreducible compact curves C_j. Exactly one of the following occurs.*

1. *If C' is a non-zero divisor supported on C then $(C', C') < 0$.*
2. *There exists a divisor $D_0 = \sum \alpha_j C_j$ such that $\alpha_j > 0$ for all j, $(D_0, D_0) = 0$, and for any effective divisor D supported on C we have $(D, D) \leq 0$ with equality if and only if D is a multiple of D_0.*
3. *There exist integers $\alpha_j > 0$ such that the divisor $D' = \sum \alpha_j C_j$ is positive and $(D', C_j) > 0$ for all j.*

Note that (*3*) in Lemma 3.3 is the same as condition $(\star)$ in Theorem 3.2. To prove Lemma 3.3, one can rephrase it as a statement about indecomposable quadratic forms that was first proved by Vinberg [29] (note that Vinberg uses a form with negative off-diagonal entries, which reverses the roles of positive and negative vectors). We make some comments about how to deduce Lemma 3.3 from [29]. That the restriction of the intersection form to the subspace of $H^2(S; \mathbb{Z})$ is indecomposable is a direct consequence of the connectedness of the reduced curve C. Since the intersection matrix is real and symmetric, it is diagonalizable over $\mathbb{R}$ with real eigenvalues. The matrix is nonsingular (resp. has rank one less than its size) if and only if all its eigenvalues (resp. all but one of its eigenvalues) are nonzero. The Lemma is now clear from combining Theorem 3 and Proposition 12 of [29].

We now state the existence theorem for positive effective Fuchsian divisors that will complete the proof of Theorem 1.2.

Theorem 3.4. *Let S be a smooth compact irreducible arithmetic quotient of $\mathbf{H}^2 \times \mathbf{H}^2$ or $\mathbf{H}^2_{\mathbb{C}}$, and suppose that S contains a Fuchsian curve. Then there is a connected reduced Fuchsian divisor D on S such that D precisely* *supports a positive effective Fuchsian divisor D' on S for which condition $(\star)$ of Theorem 3.2 holds.*

Proof. Note that under the assumptions of the theorem, S contains infinitely many commensurability classes of Fuchsian curves by Proposition 2.1 and Theorem 2.2. Since $H_2(S, \mathbb{Z})$ is finitely generated and S contains infinitely many distinct Fuchsian curves, we conclude that there is a linear dependence between homology classes of Fuchsian curves, and thus are effective Fuchsian divisors D_1 and D_2 on S such that D_1 and D_2 share no common irreducible component, D_1 is not the zero divisor, and we have an equality $[D_1] = [D_2]$ in $H_2(S, \mathbb{Z})$. It follows from invariance of the intersection pairing under homological equivalence that

$$(D_1, D_1) = (D_1, D_2) \geq 0.$$

Since (D_1, D_1) is the sum of the self intersections of the connected components of D_1, there must be a connected component D_1' of D_1 that is an effective Fuchsian divisor with $(D_1', D_1') \geq 0$.

Let $D = \sum C_j$ be the connected reduced Fuchsian divisor determined by D_1'. If $(D_1', D_1') > 0$, we are done by (*3*) in Lemma 3.3 by taking $D_1' = D'$. If $(D_1', D_1') = 0$, we claim that there exists a Fuchsian curve C_0 on S such that:

1. C_0 is not one of the irreducible components of D;
2. $(D, C_0) > 0$.

Assuming the claim, there is an integer $N > 0$ such that $D' = C_0 + N D_1'$ is a connected effective Fuchsian divisor precisely supported on $C_0 + D$ with $(D', D') > 0$, which completes the proof of the theorem as argued above.

To prove the claim, let C_0' be any Fuchsian curve on S such that C_0' is not commensurable with any Fuchsian curve in the support of D. Such a curve exists since S contains infinitely many distinct commensurability classes of Fuchsian curves. We will prove that there is a curve C_0 on S commensurable with C_0' such that $(D, C_0) > 0$.

Fix any irreducible component C_1 of the support of D, let Σ_1 be a representative for the conjugacy class of arithmetic Fuchsian subgroups of Γ associated with C_1 and Σ_0' a choice of Fuchsian subgroup associated with C_0'. Let $F_j : \mathbf{H}^2 \to X$, $j = 1, 0$, be the holomorphic totally geodesic embedding of universal coverings associated with the choice of Σ_1 and Σ_0', respectively.

Consider the *commensurator* $\mathrm{Comm}(\Gamma)$ of Γ in $G = \mathrm{Isom}(X)$, that is, the subgroup consisting of those $g \in G$ such that Γ and $g \Gamma g^{-1}$ are commensurable. By [3], $\mathrm{Comm}(\Gamma)$ is a dense subgroup in the analytic topology on G.

Let ϕ_0 be any element of G so that $\phi_0(F_0(\mathbf{H}^2))$ intersects $F_1(\mathbf{H}^2)$ transversally in a single point. By density, we can approximate ϕ_0 by $\phi \in \mathrm{Comm}(\Gamma)$ so that $\phi(F_0(\mathbf{H}^2))$ satisfies the same condition. Then $\phi\Gamma\phi^{-1}$ is a lattice in G with Fuchsian subgroup $\phi\Sigma_0'\phi^{-1}$ stabilizing $\phi(F_0(\mathbf{H}^2))$.

Since ϕ commensurates Γ, $\Sigma_\phi = \Gamma \cap (\phi\Sigma_0'\phi^{-1})$ is a Fuchsian subgroup of Γ stabilizing $\phi(F_0(\mathbf{H}^2))$. Let $\Sigma_0 \supseteq \Sigma_\phi$ be the maximal Fuchsian subgroup of Γ stabilizing $\phi(F_0(\mathbf{H}^2))$. This determines a Fuchsian curve C_0 on S commensurable with C_0' and, since $\phi(F_0(\mathbf{H}^2))$ intersects $F_1(\mathbf{H}^2)$ nontrivially, C_0 intersects C_1 nontrivially and transversely in S.

If $C_0 = C_j$, for some C_j supported on D, then C_0' and C_j would be commensurable, since C_0 and C_0' are commensurable. Since our assumptions imply that C_0 is not commensurable with any C_j, we see that C_0 is not an irreducible component of C and that $C_0 + D$ is a reduced Fuchsian curve on S that precisely supports the positive effective Fuchsian divisor $D' = C_0 + ND_1'$, for N chosen as above. This completes the proof of the claim, and thus the proof of the theorem. $\square$

Proof of Theorem 1.2. Let S satisfy the conditions of Theorem 1.2. Then Theorem 3.4 applies, so there is a Fuchsian divisor D on S satisfying the conditions of Theorem 3.2. Theorem 1.2 follows immediately. $\square$

Remark. The existence of positive effective Fuchsian divisors would, of course, be trivial if embedded Fuchsian curves have positive self-intersection. However, an embedded Fuchsian curve of genus g on a smooth compact complex hyperbolic surface has self-intersection $(1-g)$ [10, Corollary 2.3]. One can similarly show in the $\mathbf{H}^2 \times \mathbf{H}^2$ setting that the self-intersection is $2(1-g)$.

Remark. We now note that one can give upper bounds for the number of Fuchsian curves that appear in Theorem 1.2. Key in the proof is finding a linear dependence between the homology classes of Fuchsian curves inside $H_2(S;\mathbb{Z})$. Since Fuchsian curves lie in $H^{1,1}(S)$, an upper bound for the number of curves is $\dim_{\mathbb{R}} H^{1,1}(S) + 1$. Using Hirzebruch Proportionality and Noether's Theorem, one can show that

$$\frac{1}{2}e(S) + 1 \quad (X = \mathbf{H}^2 \times \mathbf{H}^2)$$

$$\frac{1}{3}e(S) + 2\dim_{\mathbb{C}}(\mathrm{Alb}(S)) + 1 \qquad (X = \mathbf{H}^2_{\mathbb{C}})$$

curves suffice, where $e(S)$ is the topological Euler characteristic and $\mathrm{Alb}(S)$ the Albanese variety (see §4). Since Euler characteristic is proportional to volume, one can express $e(S)$ in terms of the algebraic group associated with the surface S via volume formulas for arithmetic lattices. See [3] and [24] for precise descriptions of these formulas for $(\mathbf{H}^2)^n$ and $\mathbf{H}^2_{\mathbb{C}}$, respectively.

4 Albanese varieties of arithmetic complex hyperbolic 2-manifolds

In this section, let $S = \Gamma\backslash X$ be the quotient of $X = \mathbf{H}^2_{\mathbb{C}}$ by a torsion free cocompact arithmetic lattice of simple type. The Albanese variety of S is the minimal abelian variety $\mathrm{Alb}(S)$ such that any holomorphic mapping of S to an abelian variety factors through a map to $\mathrm{Alb}(S)$ (see [11, p. 331]). It was first shown by Kazhdan [14] that S always has a finite sheeted covering S' with nonzero first betti number, which occurs if and only if S' has a nontrivial Albanese. The Albanese variety of a complex projective algebraic curve C is its Jacobian variety $Jac(C)$.

Proof of Theorem 1.1. Let Γ be as in the statement of the theorem, and let $f : S \to A$ be the projection of $S = \Gamma\backslash\mathbf{H}^2_{\mathbb{C}}$ onto a simple factor A of $\mathrm{Alb}(S)$. By [11, p. 331] this induces a surjective homomorphism

$$r : \Gamma \to \mathbb{Z}^n = H_1(A, \mathbb{Z}),$$

where $n = \dim_{\mathbb{R}}(A)$.

Let $C_1, \dots, C_r$ and $D = \sum \alpha_j C_j$ be as in Theorem 1.2. The image Γ' of $\pi_1(|D|)$ in Γ has finite index in Γ. Consider the induced continuous mapping from $|D|$ to the simple abelian variety A. The image of the induced homomorphism $\pi_1(|D|) \to \mathbb{Z}^n$ has finite index in $\mathbb{Z}^n$ since Γ' has finite index in Γ and $r : \Gamma \to \mathbb{Z}^n$ is surjective.

Suppose that there an irreducible component C_j of D whose image under $f : S \to A$ is not a point. Then $f(C_j)$ is a curve, and the map from the normalization $C_j^{\#}$ of C_j to $f(C_j)$ is finite and factors through the Abel–Jacobi map from $C_j^{\#}$ to $Jac(C_j^{\#})$. Therefore $Jac(C_j^{\#})$ has nonzero image in A and, since A is simple, A is a quotient of $Jac(C_j^{\#})$.

Thus we can now suppose that the image of each C_j under $f : S \to A$ is a point. Since D is connected, it follows that $f(|D|)$ is a point on A. However, this implies that a sufficiently small neighborhood of $|D|$ maps to a contractible neighborhood of a point on A. It follows immediately that $\pi_1(|D|)$ has trivial image in $\mathbb{Z}^n$, which is a contradiction. This proves the theorem. $\square$

It would be interesting to know the following converse (cf. Corollaire 6 in the Introduction to [2]).

Question. Let S be a compact complex hyperbolic surface and C a Fuchsian curve on S. Does there exist a finite étale covering S' of S such that $Jac(C^{\#})$ is isogenous to a factor of $\mathrm{Alb}(S)$?

When S is arithmetic of simple type and $\pi_1(S)$ is a congruence arithmetic lattice, $\mathrm{Alb}(S)$ has complex multiplication [20]. However, one can construct Fuchsian curves C on certain congruence surfaces S such that $Jac(C^{\#})$ is not CM, i.e., there exist congruence arithmetic Fuchsian groups such that the Jacobian does not have complex multiplication. Therefore, even when S and C are congruence, a positive answer to the above question will necessarily require a better understanding of noncongruence coverings.

Remark. It was originally shown by Matsushima–Shimura [18] that irreducible quotients of $\mathbf{H}^2 \times \mathbf{H}^2$ have trivial first cohomology over $\mathbb{C}$, and thus have trivial Albanese variety. In particular, the results in this section are trivial for irreducible lattices in $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$. For reducible quotients, the Albanese is easily seen to be isogenous to a product of Jacobians.

References

[1] W. L. Baily, Jr. On compactifications of orbit spaces of arithmetic discontinuous groups acting on bounded symmetric domains. In *Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965)*, pages 281–295. Amer. Math. Soc., 1966.

[2] N. Bergeron and L. Clozel. Spectre automorphe des variétés hyperboliques et applications topologiques. *Astérisque*, (303), 2005.

[3] A. Borel. Density and maximality of arithmetic subgroups. *J. Reine Angew. Math.*, 224:78–89, 1966.

[4] A. Borel. Commensurability classes and volumes of hyperbolic 3-manifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 8(1):1–33, 1981.

[5] F. Campana. Negativity of compact curves in infinite covers of projective surfaces. *J. Algebraic Geom.*, 7(4):673–693, 1998.

[6] T. Chinburg, E. Hamilton, D. D. Long, and A. W. Reid. Geodesics and commensurability classes of arithmetic hyperbolic 3-manifolds. *Duke Math. J.*, 145(1):25–44, 2008.

[7] L. Clozel and T. N. Venkataramana. Restriction of the holomorphic cohomology of a Shimura variety to a smaller Shimura variety. *Duke Math. J.*, 95(1):51–106, 1998.

[8] S. S. Gelbart and J. D. Rogawski. L-functions and Fourier-Jacobi coefficients for the unitary group U(3). *Invent. Math.*, 105(3):445–472, 1991.

[9] W. M. Goldman. *Complex hyperbolic geometry.* Oxford Mathematical Monographs. 1999. Oxford Science Publications.

[10] William M. Goldman, Michael Kapovich, and Bernhard Leeb. Complex hyperbolic manifolds homotopy equivalent to a Riemann surface. *Comm. Anal. Geom.*, 9(1):61–95, 2001.

[11] P. Griffiths and J. Harris. *Principles of algebraic geometry.* Wiley Classics Library. John Wiley & Sons Inc., 1994.

[12] Robin Hartshorne. *Algebraic geometry.* Springer-Verlag, 1977. Graduate Texts in Mathematics, No. 52.

[13] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 80. Academic Press Inc., New York, 1978.

[14] D. Kazhdan. Some applications of the Weil representation. *J. Analyse Mat.*, 32:235–248, 1977.

[15] R. Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete.* Springer-Verlag, Berlin, 2004.

[16] C. Maclachlan and A. W. Reid. *The arithmetic of hyperbolic 3-manifolds*, volume 219 of *Graduate Texts in Mathematics.* Springer-Verlag, 2003.

[17] G. A. Margulis. *Discrete subgroups of semisimple Lie groups*, volume 17 of *Ergebnisse der Mathematik und ihrer Grenzgebiete.* Springer-Verlag, 1991.

[18] Yozô Matsushima and Goro Shimura. On the cohomology groups attached to certain vector valued differential forms on the product of the upper half planes. *Ann. of Math. (2)*, 78:417–449, 1963.

[19] D. B. McReynolds and A. W. Reid. The genus spectrum of a hyperbolic 3-manifold. preprint, 2009.

[20] V. K. Murty and D. Ramakrishnan. The Albanese of unitary Shimura varieties. In *The zeta functions of Picard modular surfaces*, pages 445–464. Univ. Montréal, 1992.

[21] T. Napier and M. Ramachandran. The L^2 $\overline{\partial}$-method, weak Lefschetz theorems, and the topology of Kähler manifolds. *J. Amer. Math. Soc.*, 11(2):375–396, 1998.

[22] M. V. Nori. Zariski's conjecture and related problems. *Ann. Sci. École Norm. Sup. (4)*, 16(2):305–344, 1983.

[23] G. Prasad and A. S. Rapinchuk. Weakly commensurable arithmetic groups and isospectral locally symmetric spaces. *Publ. Math. Inst. Hautes Études Sci.*, (109):113–184, 2009.

[24] G. Prasad and S.-K. Yeung. Fake projective planes. *Invent. Math.*, 168(2):321–370, 2007.

[25] A. W. Reid. Isospectrality and commensurability of arithmetic hyperbolic 2- and 3-manifolds. *Duke Math. J.*, 65(2):215–228, 1992.

[26] W. Scharlau. *Quadratic and Hermitian forms*, volume 270 of *Grundlehren der Mathematischen Wissenschaften.* Springer-Verlag, 1985.

[27] J. Schwermer. Geometric cycles, arithmetic groups and their cohomology. *Bull. Amer. Math. Soc. (N.S.)*, 47(2):187–279, 2010.

[28] M.-F. Vignéras. *Arithmétique des algèbres de quaternions*, volume 800 of *Lecture Notes in Mathematics.* Springer, 1980.

[29] È. B. Vinberg. Discrete linear groups that are generated by reflections. *Izv. Akad. Nauk SSSR Ser. Mat.*, 35:1072–1112, 1971.

[30] E. B. Vinberg, editor. *Lie groups and Lie algebras. II*, volume 21 of *Encyclopaedia of Mathematical Sciences.* Springer-Verlag, Berlin, 2000.